

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MaxCel[®]

PLANT GROWTH REGULATOR SOLUTION

ACTIVE CONSTITUENT: 19 g/L 6-BENZYLADENINE

For reducing fruit set and increasing fruit size and improving firmness of harvested apples as per the DIRECTIONS FOR USE table.

GENERAL INSTRUCTIONS

MaxCel is a secondary thinner designed for use on trees treated with a primary thinner (such as NAA) at or about flowering. Best results from MaxCel will be obtained when foliage is evenly and thoroughly covered with the spray.

Timing and temperature at time of application is critical. MaxCel does not increase the incidence of russet when applied alone at label rates. An increase in vegetative growth may be observed following application of MaxCel.

MIXING

MaxCel mixes easily with water. Add the required amount to a clean spray tank about half filled with water. Agitate while bringing the total volume of the water to the required level. Use the spray solution within 24 hours after mixing. Addition of a non-ionic wetting agent may improve coverage.

APPLICATION

Apply in sufficient water to achieve thorough and even coverage of leaves. Coverage of lower leaf surfaces is important.

Apply when king fruitlets are between 7-10 mm in diameter. This normally occurs 10 to 25 days after full bloom.

Spray on a day when the daytime temperature is increasing to a predicted daily maximum of greater than 15°C.

COMPATIBILITY

It is recommended that MaxCel not be mixed with other pesticides, growth regulators or fertilisers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with chemical or used container.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a well ventilated area, as cool as possible. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing product for use wear cotton overalls button to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls button to the neck and wrist and washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone: 13 11 26)

SDS

Additional information is listed in the Safety Data Sheet (SDS).

IMPORTANT NOTICE

These goods are to be used only for the purpose and as specified on the label, and are not suitable for any other purpose. To the fullest extent permitted by law, we do not accept or bear any liability on any basis for any loss, damage, cost or expense, arising in any way, directly or indirectly, in connection with the goods.

® Registered Trademark of Valent BioSciences Corporation, Libertyville, Illinois, USA
© Valent BioSciences 2011

APVMA Approval No.: 61799/55873

THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF DANGEROUS GOODS BY ROAD OR RAIL

**In a Transport
Emergency
Dial 000
Police or
Fire Brigade**

**SPECIALIST ADVICE
IN EMERGENCY ONLY
ALL HOURS
AUSTRALIA WIDE
1800 024 973**

DIRECTIONS FOR USE

Restrains:

DO NOT apply when maximum daily temperature is likely to be less than 15°C.

DO NOT apply later than 30 days after full bloom.

CROP	RATE	CRITICAL COMMENTS
Apple varieties: Fuji, Gala, Golden Delicious, Pink Lady, Red Delicious Sundowner	9 L/ha in a minimum of 1200 L/ha	Use as a secondary thinner after application of a primary thinner (such as NAA) at bloom. Increase water volume as canopy density increases. Apply when king fruitlets are between 7-10 mm in diameter. This normally occurs 10 to 25 days after full bloom. Spray on a day when the daytime temperature is increasing to a predicted daily maximum of greater than 15°C.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

NOT REQUIRED WHEN USED AS DIRECTED.