

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Gro-Chem NAA 20

FRUIT THINNING AND SETTING COMPOUND.

ACTIVE CONSTITUENT: 20 g/L N.A.A. present as the sodium salt

**To assist in the prevention of pre-harvest drop of Apples and Pears.  
May also be used as a thinning spray for Apples and for striking hard and softwood cuttings.**

## GENERAL INSTRUCTIONS

When spraying for the prevention of pre-harvest fruit drop, do not allow fruit to remain on trees beyond normal harvest date otherwise fruit may become over mature.

Use a suitable wetter such as Polysorbate 20 at 125 mL/100 L of spray volume.

NAA must be absorbed into the tree to be effective. Therefore slow drying conditions after spraying, high volume spraying and the use of wetting agent will enhance effectiveness.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to re-cycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. After use and before eating, drinking and smoking, wash hands, arms and face thoroughly with soap and water. When preparing spray wear face shield or goggles.

Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS).

## IMPORTANT NOTICE

These goods are to be used only for the purpose and as specified on the label, and are not suitable for any other purpose. To the fullest extent permitted by law, we do not accept or bear any liability on any basis for any loss, damage, cost or expense, arising in any way, directly or indirectly, in connection with the goods.

APVMA Approval No.: 60577/1L-20L/0107

In a Transport Emergency  
**Dial 000**  
Police or Fire Brigade

**SPECIALIST ADVICE  
IN EMERGENCY ONLY  
1800 024 973  
ALL HOURS - AUSTRALIA WIDE**

## DIRECTIONS FOR USE

CROP	ACTION	RATE	CRITICAL COMMENTS
Apples and Pearsars	Assist prevention of pre-harvest fruit drop	1 L / 2000 L of water (10 ppm) 500 mL / 2000 L of water (5 ppm)	Apply as a high volume spray 12 to 14 days before harvest to thoroughly wet fruit and foliage. A second, half-strength spray (5 ppm) may be necessary if conditions favour excessive fruit drop.
Apples	Fruit thinning	250 to 1000 mL per 2000 L of water (2:5 – 10 ppm)	Apply as a high volume spray 10 to 14 days after full bloom to the top two-thirds of the tree for uniform thinning. For specific varieties, consult local Dept. of Agriculture. In Tasmania, apply, particularly to Golden Delicious, from balloon blossom to 7 days after full bloom.
Hard or Softwood cuttings	Root striking	12.5 mL to 50 mL per 10 L of water.	Dip base of cutting to depth of 5 mm into NAA solution for approximately 5 seconds. Allow to dry and insert half the length of the cutting into striking medium. Use the stronger solution for hardwood cuttings, weaker solution for softwood cuttings.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD – DO NOT APPLY LATER THAN 1 DAY BEFORE HARVEST.**